


RESEARCH ARTICLE

Surveying and mapping cereals and legumes wild relatives in Mount Hermon (Bekaa, Lebanon)

Eliane Sayde¹ | Lamis Chalak¹ | Safaa Baydoun² | Ali Shehadeh³ | Hicham El Zein⁴  | Jostelle Al Beyrouthy³  | Mariana Yazbek³ 

¹Faculty of Agronomy, Department of Plant Production, Lebanese University, Beirut, Lebanon

²Research Center for Environment and Development, Beirut Arab University, Bekaa, Lebanon

³International Center for Agricultural Research in Dry Areas (ICARDA), Beirut, Lebanon

⁴Independent Researchers, Beirut, Lebanon

Correspondence

Mariana Yazbek, International Center for Agricultural Research in Dry Areas (ICARDA), Beirut, Lebanon.
Email: m.yazbek@cgiar.org

Abstract

Crop Wild Relatives (CWR) should be highly prioritized, monitored, and conserved as they have an immense effect on sustainability and livelihood. In this study we aim to survey and map cereal and legume wild relatives of Fabaceae and Poaceae families. Mount Hermon, Bekaa side, Lebanon. A set of 46 CWR species were targeted based on desk selection analysis and prioritization by the International Center for Agricultural Research in Dry Areas genebank for their potential importance in breeding programs. A botanical survey of 17 sites of the various habitats of Mount Hermon was performed during April–June 2021 using a systematic transect/quadrat sampling method. Recorded genera and species were accurately georeferenced and then mapped with the DIVA-GIS program. In total, 854 occurrences were observed belonging to 34 species of Fabaceae and 12 species of Poaceae. High H' Shannon diversity values were recorded in three sites (Al Fakiaa, Sham El Hafour and Ain Ata- al Berke) of the Mount with values ranking between 2.45 and 2.83. This was confirmed by the richness distribution maps of genera and species. Richness distribution maps provide relevant clues on candidate sites for high concentrations of each of the species under study. At least the three sites, hosting 87% of the surveyed CWR's species, can be considered for further in situ conservation actions.

KEYWORDS

botanical survey, cereals and legumes, CWR, distribution mapping, DIVA-GIS, mount Hermon

TAXONOMY CLASSIFICATION

Biodiversity ecology, Conservation ecology

1 | INTRODUCTION

Crop Wild Relatives (CWR) are a valuable source of useful diversity that can improve crop performance to adapt to ever-changing conditions and to overcome the constraints caused by pests, diseases, and abiotic stresses (Bohra et al., 2021, Perrino & Wagensommer, 2021). They represent an extended gene pool that

can be tapped for the breeding of novel varieties with adaptability traits to various environmental stresses (Maxted et al., 2016). CWR are also essential for sustainable agriculture and to maintain agrobiodiversity in a scenario of climate change and its unpredictable outcomes (Bierschenk et al., 2020). Nevertheless, many CWR genetic resources are threatened in their natural habitats and/or not fairly represented in genebank and conservation initiatives

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2024 The Authors. *Ecology and Evolution* published by John Wiley & Sons Ltd.

(Bohra et al., 2021; Castañeda-Álvarez et al., 2016; Ceccarelli et al., 2012; Kell et al., 2012). These conservation gaps limit therefore the range of useful plant diversity that should be available for present and future generations (Khoury et al., 2019). These CWR should be considered as a priority for conservation, monitoring, and sustainable utilization.

Technology-based approaches such as generating distribution maps are deployed as very important tools to illustrate the current richness of CWR, predicted future richness, and its changing trends (Jarvis et al., 2008). Such maps are developed as a valuable tool for guiding the implementation of appropriate conservation strategies. More specifically, they can be used for prioritizing the hotspot areas for in situ conservation such as the establishment of CWR reserves (Ratnayake et al., 2021). For instance, mapping the richness and distribution of prioritized CWR in Zambia helped identify sites for active in situ conservation as a part of a national plan for the development of conservation and sustainable use of CWR (Ng'uni et al., 2019).

Located at the Eastern shore of the Mediterranean Basin, Lebanon represents one of the most important agrobiodiversity hotspots, not only for wild species diversity but also for cultivated plant diversity (Pacocco et al., 2018). Lebanon is also well recognized for its wealth in CWR (Nail, 2019), with a large share of Poaceae and Fabaceae families (Chalak et al., 2020; Zair et al., 2018). The country is also part of the Fertile Crescent that is known to have been the cradle of agriculture as having a very rich and unique biodiversity of flora owed to its mountainous topography and the great diversity of its climatic conditions (Chalak et al., 2011, 2020). It has a high percentage (12%) of endemic plant species which are mostly located on the high summits of the two mountain ranges spanning parallel to the sea coast (i.e., Lebanon Mountains and Anti-Lebanon Mountains) (Chalak et al., 2020).

Mount Hermon, also known as Jabal al-Shaykh, is a mountain cluster constituting the southern end of the Anti-Lebanon Mountains range (Baydoun et al., 2015). The mountain offers a wide range of ecosystem products mainly in terms of fresh water and agricultural crops. Mount Hermon was described to be a high mountain plateau characterized by the rarity, richness, and uniqueness of vascular plants. It was identified as an Important Plant Area in Lebanon where endemic and/or threatened species are found (Bou Dagher Kharrat et al., 2018; IUCN, 2001) and where more than hundred species specific to Mount Hermon and the Anti-Lebanon range have been counted (Arnold et al., 2015; Medail & Quezel, 1997). A Richness Index (i.e., a numerical score assigned to each cell based on the number of species present, using a scale ranging from 0 to 5 and calculated using the Jenks natural breaks classification method (1967)) scored 4 referring to 259 species harbored by the Mount was reported (Bou Dagher Kharrat et al., 2018). Many of these species are characteristic of the semi-arid Eastern Mediterranean climate and are endemic to Lebanon and Syria (Arnold et al., 2015; Bou Dagher Kharrat et al., 2018). The natural wealth of Mount Hermon has been subject to pressures impacting its plants and all other types of wildlife. Habitat

fragmentation, overexploitation of resources, grazing, invasive alien species, urban expansion, and increasing impact of climate change are among the challenges facing the mountain and its surrounding communities (Arnold et al., 2015).

Recently and within its mission of collecting, conserving, and utilizing dryland agrobiodiversity, the International Center for Agricultural Research in Dry Areas (ICARDA) took the initiative to survey and inventory the CWR growing in Lebanon, and more particularly in the Bekaa province to bridge the gaps of ex situ conservation (Amri et al., 2016; Nail, 2019). In this study, we aim to survey and map CWR in the mountain with an emphasis on Fabaceae and Poaceae species particularly selected for their genetic and economic importance. The study will help bridging the CWR gaps of ICARDA genebank for potential use in breeding programs. It will also provide baseline knowledge to identify hotspots for CWR in situ conservation and management.

2 | MATERIALS AND METHODS

2.1 | Study area

This study was conducted on the western slopes of Mount Hermon, Bekaa side, covering an area of about 190 km² that extends between 1067 and 2800 m of altitude. The study area was delimited using a tool in DIVAGIS which is set to layer an ecogeographic grid dividing the study area into georeferenced cells (48 cells) of 2 × 2 km (i.e., 4 km² quadrants), covering the entire area. Seventeen cells out of the total 48 cells representing the different habitats described by Chalak et al., n.d. and capturing the heterogeneity of plant communities of the Mount while considering accessibility were selected based on their accessibility on Mount Hermon (Table 1). The location of Mount Hermon study area in Lebanon is visualized in Figure S1.

2.2 | CWR targeted species

A desk analysis of various published sources including Vincent et al. (2013) global priority list of CWR, [Global Biodiversity Information Facilities (GBIF, 2021) (<http://www.gbif.org>)], Genesys (<https://www.genesys-pgr.org/>), Euro Med Plant Base (<http://www.emplantbase.org>), the New Flora of Lebanon and Syria (Mouterde 1966, 1970; Mouterde, 1986) and Lebanon-Flora (<http://www.lebanon-flora.org>), and a dissertation (Al-Atawneh et al., 2008); and literature related to the flora in Mount Hermon (Baydoun et al., 2015) was first conducted as a pre-selection step of CWRs found in the Mount. Following this step, 15 CWRs genera belonging to Fabaceae and Poaceae which are two major families present in Lebanon were selected for this study based on their genetic and economic values according to their importance in ICARDA's gene bank which is the main ex situ conservation center for cereals, food legumes, forage, and rangeland species in the country, as

TABLE 1 Surveyed areas in Mount Hermon with their respective latitude, longitude, altitude, and general dominating habitats.

Study sites	Latitude	Longitude	Altitude (m)	Habitat (Chalak et al. (n.d.))
Tanoura- Al Jouk	33.46647	35.80224	1104	Woodland— <i>Quercus coccifera</i> Rocky grassland
Arid El Dalia	33.469006	35.8216	1120	Woodland— <i>Quercus coccifera</i>
Al Fakiaa	33.45338	35.8132	1135	Woodland— <i>Quercus coccifera</i>
Sahel Ayha-Wadi Al Byara	33.50786	35.8668	1173	Agriculture
Wadi El Raheb	33.44057	35.81472	1219	Woodland— <i>Quercus infectoria</i>
Sham El Hafour	33.430588	35.80259	1278	Agriculture Woodland— <i>Quercus infectoria</i>
Hima El Kadirin	33.447124	35.822078	12 79	Woodland— <i>Quercus coccifera</i> Woodland— <i>Quercus infectoria</i>
Al Yebse- Wadi El Botem	33.47244	35.846858	1317	Woodland and grassland— <i>Quercus infectoria</i> Woodland— <i>Quercus infectoria</i>
Ain Ata- Marj El Tout	33.488559	35.88283	1335	Thickets of <i>Quercus</i> and <i>Crataegus</i> Agriculture
Ain Ata-Wadi Ain El Sayegh	33.41904	35.80208	1373	Thickets of <i>Quercus</i> and <i>Crataegus</i>
Ain Ata-Jabal El Khan	33.41807	35.78079	1404	Woodland— <i>Quercus infectoria</i>
Rashaya El Wadi-Wadi Samouk	33.472748	35.866075	1409	Thickets of <i>Quercus</i> and <i>Crataegus</i>
Ain Ata- Al Berke	33.432384	35.782907	1409	Agriculture Woodland— <i>Quercus coccifera</i>
Rouwayset El Rafiaa-Sammouk	33.462342	35.8852	1554	Thickets of <i>Quercus</i> and <i>Crataegus</i>
Chebaa-Ouyoun Jena	33.3909	35.791365	1575	Woodland— <i>Prunus dulcis</i> — <i>Quercus look</i>
Jourit Al Nkar-Al Chok	33.448598	35.845698	1637	Thickets of <i>Quercus</i> and <i>Crataegus</i> Woodland— <i>Prunus dulcis</i> — <i>Quercus look</i>
Fahet Jernaya- Beb El Hawa	33.425322	35.839478	2254	Oro-Mediterranean shrublands, rocks and screes

Note: Latitude and longitude correspond to the center of the cells.

well as continuously promoting the utilization of these wild species in experiment targeting world food security. Over these 15 CWRs genera, eight represented Fabaceae consisting of *Lathyrus*, *Lens*, *Medicago*, *Pisum*, *Trifolium*, *Trigonella*, *Coronilla*, and *Vicia*, and the remaining seven genera belonged to Poaceae including *Aegilops*, *Avena*, *Bromus*, *Hordeum*, *Lolium*, *Poa*, and *Triticum*.

2.3 | Botanical survey

Ten field botanical survey missions were conducted in the natural habitats of the mountain between April and June 2021, exploring the 17 sites, assisted by a guide from the local habitants. Transect/quadrates survey method was used to assess the CWR status. We took into consideration the cells that represent the various types of habitats, noting that some cells designate more than one habitat type. Consequently, each designated cell had to include two transects positioned in a manner to adequately represent the plant communities growing in these habitats of Mount Hermon. Each transect was around 100m trajectory and comprised five quadrates of 1m² with an interval distance of 25m. A total of 170 quadrates were surveyed. Data related to the site coordinates, general geomorphology, general habitat, land cover, and slope were collected.

For the CWR species, records of their Richness, Occurrence, Frequency, and Density were collected, with species nomenclature based on Mouterde (1966, 1970, Mouterde, 1986), the online Euro Med Plant Base (2006) (<http://www.emplantbase.org>), and the help of specialized botanists.

2.3.1 | Species indices

Indices related to species considered in this study were: (i) *Species occurrence* considered to be the species presence/absence in the assessed 170 quadrates; (ii) *Species density* defined as the number of individuals of a given species that occurs within a quadrate (MacKenzie et al., 2018); (iii) *Species frequency* calculated as the percentage of occurrences in the 170 quadrates assessed (Good, 1953).

2.3.2 | Diversity indices per site

(i) *Genus richness* and *species richness* being a useful measure of taxonomic diversity (Gaston, 1996) and calculated as number of different genera or species per site (representing the 17 cells); (ii) *Shannon diversity index (H')* that allows to quantify the

species diversity per area unit (cell) using the following formula (Magurran, 1998; Rajan, 2001):

$$H' = - \sum (P_i * \ln P_i).$$

where P_i is the proportion (n/N) of individuals of one particular species found (n) divided by (N) total number of individuals found, $\ln P_i$ is the natural log of P_i , and \sum is the sum of P_i calculations. Species with individual count below 50 were registered as densities (i.e., number of individuals of a species within a quadrat considering that quadrats are representative of the whole cell), while species with individual count equal or greater than 50 were taken as 50 for the sake of the calculation.

3 | CWR MAPPING

Data collected in botanical surveys of targeted CWR were mapped based on calculated indices described above.

3.1 | Genus and species mapping

DIVA-GIS version 7.1.7 was used to map each CWR genera and species distribution in Mount Hermon using data entry tables of identified genera and species with their nomenclature and respective accurate georeferenced occurrence locations collected during the botanical surveys, alongside their densities in quadrats.

4 | RESULTS

4.1 | Species occurrence, frequency, and density of targeted CWRs

In this study, 859 georeferenced observations in Mount Hermon were registered for 46 CWR species belonging to the 15 targeted genera, with 12 Poaceae species and 34 Fabaceae species. The most represented genus is *Trifolium* with 17 species, followed by *Medicago* with eight species, while genera like *Bromus*, *Poa*, *Coronilla*, and *Pisum* are represented with only one species (Table 2).

Trifolium stellatum L. had the highest frequency in quadrats of 61.18% with 104 occurrences, followed by *Avena sterilis* L. (58.24% frequency and 99 occurrences), *T.resupinatum* var. *resupinatum* (54.12% frequency and 92 occurrences). Oppositely, nine species recorded only one occurrence each, designating 0.59% on the frequency chart, such as *M.minima* (L.) Bart., *M.sativa* L., *T.fragiferum* L., *T.spumosum* L. and *V.sativa* L. The majority of Poaceae species registered high frequency records between 12.35% and 58.24%, with the *Hordeum* species scoring different occurrence frequencies, *H.murinum* subsp. *glaucom* (Steud.) Tzvelev (12.35%), *H.bulbosum* L. (18.82%), and the highest occurrence frequency for this genus was *H.spontaneum* C. Koch (20.59%). As for Fabaceae species, only few recorded a relatively high frequency value mainly for species

belonging to the *Trifolium* genus, *T.pilulare* Boiss. (22.94%) and *T.scutatum* Boiss. (34.12%). Species densities per quadrat fluctuated for the same species depending on the habitat types. All the Poaceae species were present in high densities within the quadrats, as most values were in the range of (1–50), except for *Poa* (5–8) and *Triticum* (0–1) genera that had low species density. For Fabaceae, density ranges were highly fluctuating within the same genera such as *T.boissieri* Guss. scores (1–50) and *T.subterraneum* var. *brachycalycinum* Katzn. & Morley which counted only one time in a quadrat.

4.2 | Diversity indices of targeted CWRs

The surveyed sites varied in their genus and species richness (Table 3). Up to 35 species belonging to 12 genera were found in Sham El Hafour, 36 species belonging to 9 genera in Al Fakiaa, and 24 species belonging to 9 targeted genera in Ain Ata- Al Berke. Fahet Jernaya- Beb El Hawa scored the lowest richness with 6 species belonging to 4 genera.

Shannon index representing species richness in the assessed cells was calculated to describe and compare the CWR taxonomical diversity among the visited areas of Mount Hermon. The index fell in the range of 1.23 and 2.82 with the majority of the cells demonstrating high important values, particularly in the sites of Sham El Hafour (2.83), Al Fakiaa (2.71), Ain Ata- Al Berke (2.45), and Wadi El Raheb (2.41) which were the most diversified in targeted CWR species compared to other sites in the Mount.

4.3 | CWR mapping

4.3.1 | CWR mapping based on genus richness

The sites with the highest CWR genus richness were found in Al-Fakiaa, Hima El Kadirin, and Sham El Hafour where the dominant habitats are woodland—*Quercus coccifera* and woodland—*Quercus infectoria* counting 10–11 genera in these sites (Figure 1a). At the opposite, the sites with the lowest CWR genus richness included Fahet Jernaya-Beb El Hawa characterized with an oro-Mediterranean shrublands, rocks, and screes habitat type, and the area of Marj El Tout where the dominant habitat is thickets of *Quercus* and *Crataegus* accompanied with agricultural lands. In these sites, the CWR genus richness did not exceed the range of 1–4 genera. Sites with CWR richness of 5–9 genera were found in the dominant habitats of: Woodland—*Quercus coccifera*, Woodland—*Quercus infectoria*, Woodland—*Prunus dulcis*-*Quercus look*, and agricultural lands.

4.3.2 | CWR mapping based on species richness

Figure 1b shows that the 46 identified CWR species were mainly focused in Al Fakiaa and Sham El Hafour where the dominant habitats are respectively woodland—*Quercus coccifera* and agricultural land counting 23–28 species. Contrarily, areas with the lowest richness of 1–6

TABLE 2 List of targeted CWRs in Mount Hermon, with family, frequencies (occurrences), and range of densities in quadrates where they occurred.

Species	Family	Related crop	Species frequency (%) (number of species occurrence in quadrates)	Density range in quadrates (minimum–maximum)
<i>Ae. biuncialis</i> Vis.	Poaceae	Wheat	5.29% (9)	1–50
<i>Ae. geniculata</i> Roth.	Poaceae	Wheat	4.12% (7)	1–50
<i>Ae. sp.</i>	Poaceae	Wheat	51.76% (88)	2–50
<i>Ae. triuncialis</i> L.	Poaceae	Wheat	0.59% (1)	0–17
<i>A. sterilis</i> L.	Poaceae	Oat	58.24% (99)	1–50
<i>B. sterilis</i> L.	Poaceae	Brome grass	19.41% (33)	1–50
<i>Coronilla scorpioides</i> L.	Fabaceae	Vetch	5.88% (10)	1–20
<i>H. bulbosum</i> L.	Poaceae	Barley	18.82% (32)	1–27
<i>H. murinum</i> subsp. <i>glaucum</i> (Steud.) Tzvelev	Poaceae	Barley	12.35% (21)	1–50
<i>H. spontaneum</i> C. Koch	Poaceae	Barley	20.59% (35)	1–50
<i>L. aphaca</i> L.	Fabaceae	Pea	10% (17)	1–50
<i>L. blepharicarpus</i> Boiss.	Fabaceae	Pea	7.65% (13)	1–24
<i>Lens culinaris</i> subsp. <i>orientalis</i> (Boiss.) Ponert	Fabaceae	Lentil	2.94% (5)	2–50
<i>Lolium rigidum</i> Goud.	Poaceae	Ryegrass	25.29% (43)	1–50
<i>M. coronata</i> (L.) Bart.	Fabaceae	Alfalfa	2.35% (4)	0–1
<i>M. minima</i> (L.) Bart.	Fabaceae	Alfalfa	0.59% (1)	0–1
<i>M. orbicularis</i> L.	Fabaceae	Alfalfa	2.35% (4)	2–12
<i>M. praecox</i> DC.	Fabaceae	Alfalfa	4.71% (8)	1–50
<i>M. radiata</i> L.	Fabaceae	Alfalfa	1.18% (2)	3–13
<i>M. rigidula</i> var. <i>rigidula</i> (L.) Desr.	Fabaceae	Alfalfa	2.35% (4)	1–4
<i>M. rotata</i> var. <i>rotate</i> Boiss.	Fabaceae	Alfalfa	8.82% (15)	1–17
<i>M. sativa</i> L.	Fabaceae	Alfalfa	0.59% (1)	0–50
<i>Pisum fulvum</i> Sibth. & Sm.	Fabaceae	Pea	1.18% (2)	4–6
<i>Poa pratensis</i> L.	Poaceae	Meadowgrass	2.35% (4)	5–8
<i>T. arvense</i> L.	Fabaceae	Clover	1.18% (2)	2–12
<i>T. boissieri</i> Guss.	Fabaceae	Clover	18.82% (32)	1–50
<i>T. bullatum</i> Boiss. & Hausskn. var. <i>bullatum</i>	Fabaceae	Clover	2.94% (5)	2–12
<i>T. campestre</i> Schreb.	Fabaceae	Clover	2.94% (5)	2–50
<i>T. clusii</i> Gren. & Godr var. <i>clusii</i> .	Fabaceae	Clover	1.18% (2)	0–2
<i>T. fragiferum</i> L.	Fabaceae	Clover	0.59% (1)	0–50
<i>T. glanduliferum</i> Boiss.	Fabaceae	Clover	5.88% (10)	3–50
<i>T. hirtum</i> All.	Fabaceae	Clover	1.76% (3)	4–28
<i>T. pilulare</i> Boiss.	Fabaceae	Clover	22.94% (39)	1–50
<i>T. purpureum</i> Loisel. var. <i>purpureum</i>	Fabaceae	Clover	8.82% (15)	1–50
<i>T. resupinatum</i> var. <i>resupinatum</i> L.	Fabaceae	Clover	54.12% (92)	1–50
<i>T. scabrum</i> L.	Fabaceae	Clover	2.94% (5)	1–20
<i>T. scutatum</i> Boiss.	Fabaceae	Clover	34.12% (58)	1–50
<i>T. spumosum</i> L.	Fabaceae	Clover	0.59% (1)	0–21
<i>T. stellatum</i> L.	Fabaceae	Clover	61.18% (104)	1–50
<i>T. subterraneum</i> var. <i>brachycalycinum</i> Katzn. & Morley	Fabaceae	Clover	0.59% (1)	0–1
<i>T. tomentosum</i> var. <i>glabrescens</i> (Post) Hausskn. & Bornm.	Fabaceae	Clover	5.29% (9)	1–50
<i>Trigonella spinosa</i> L.	Fabaceae	Fenugreek	0.59% (1)	0–7
<i>Triticum aestivum</i> L.	Poaceae	Wheat	0.59% (1)	0–1
<i>Vicia sativa</i> L.	Fabaceae	Vetch	0.59% (1)	0–18
<i>Vicia tenuifolia</i> Roth.	Fabaceae	Vetch	4.71% (8)	1–8

TABLE 3 Shannon diversity Index for targeted CWRs, the genus richness, and species richness in surveyed areas of Mount Hermon.

Site	Genus richness	Species richness	Shannon index
Al Fakiaa	9	36	2.71
Sham El Hafour	12	35	2.83
Wadi El Raheb	9	21	2.41
Ain Ata-Jabal El Khan	6	17	2.27
Ain Ata-Wadi Ain El Sayegh	8	11	1.64
Rashaya El Wadi-Wadi Samouk	7	18	2.03
Rouwayset El Rafiaa-Sammouk	7	12	1.998
Jourit Al Nkar-Al Chok	9	15	2.19
Al Yebse- Wadi El Botem	7	14	1.67
Hima El Kadirin	11	24	2.27
Arid El Dalia	7	16	1.58
Chebaa-Ouyoun Jena	8	15	1.23
Ain Ata- Al Berke	9	24	2.45
Ain Ata- Marj El Tout	6	14	1.89
Fahet Jernaya- Beb El Hawa	4	6	1.45
Sahel Ayha-Wadi Al Byara	6	23	1.65
Tanoura- Al Jouk	9	15	2.29

species were found in oro-Mediterranean shrublands, rocks and scree, rocky grassland, woodland—*Quercus infectoria* and thickets of *Quercus* and *Crataegus* habitat types. Ain Ata-El Berke could be considered to have a moderately high CWR species richness counting between 18 and 22 CWR species in this site characterized as agricultural land.

4.3.3 | CWR mapping based on Shannon diversity index

The obtained values of Shannon diversity index varied from one habitat to another in the same cell as represented in the distribution map generated for representing CWR species diversity across the mountain habitats (Figure 1c).

4.4 | Distribution of wild relatives of cereals

4.4.1 | *Aegilops*

Aegilops species occurred in all the 17 studied sites (Figure 2a) where *Ae. biuncialis* Vis, *Ae. geniculata* Roth., and *Ae. triuncialis* L. were found, while other occurrences were registered as *Ae. sp.* when certainty of species identification was not adequate.

4.4.2 | *Avena*

Only one species of *Avena* was found, *A. sterilis* L., with a wide distribution all over the mountain (Figure 2b) and mainly in the sites of Sahel Aiha-Wadi el Byara and in Marj El Tout.

4.4.3 | *Bromus*

This genus was found under only one species, *B. sterilis* L. (Figure 2c). Although highly occurring in sites that are identified as woodlands, Arid El Dalia and Rashaya El Wadi-Wadi Samouk.

4.4.4 | *Hordeum*

This genus is widely distributed throughout Mount Hermon and is found in all the habitats (Figure 2e). Three species were identified in different occurrence numbers simultaneously *H. murinum* subsp. *glaucom* (21), *H. bulbosum* L. (32), and *H. spontaneum* C. Koch (35) (Figure 2f). Species richness analysis shows that all three species of *Hordeum* are found in six sites, most importantly in Sahel Aiha- Wadi El Byara where all the species are found.

4.4.5 | *Lolium*

This genus occurred in all studied sites and habitats of the mountain, while only one species was identified, *Lolium rigidum* Goud. This species is mainly located in Sahel Aiha-Wadi el Byara (agricultural land) and Marj El Tout (woodland) (Figure 2d).

4.4.6 | *Poa* and *Triticum*

Both genera occurred only one time during the survey, under the species *Poa pratensis* L. and *T. aestivum* L. therefore relevant information are too limited in this study.

4.5 | Distribution of wild relatives of legumes


4.5.1 | *Lathyrus*

Distribution analysis for this genus shows its occurrence in 10 out of 17 surveyed sites (Figure 3a). *Lathyrus* species richness revealed two identified species, *Lathyrus blepharicarpus* Boiss. and *L. aphaca* L. (Figure 3b), that were both found at the same time in Al Fakiaa, Hima El Kadirin, Ain Ata el Berke characterized as woodlands, and also in Sham El Hafour and Chebaa-Ouyoun Jena both considered as grassland and shrublands.

4.5.2 | *Medicago*

Species are distributed in a very random manner all over the different habitats of the Mount where 10 cells out of 17 reported this genus (Figure 3c). Eight different *Medicago* species were identified with *Medicago rotata* var. *rotata* Boiss. having the highest occurrence number, followed by *M. preacox* DC. with an occurrence value of 15, then *M. coronata* (L.) Bart., *M. orbicularis* L., and

FIGURE 1 (a) Distribution map of CWR genus richness, (b) CWR species richness, and (c) Map representing CWR diversity ranges using Shannon index in Mount Hermon (generated by DIVA-GIS).






FIGURE 2 Distribution map of (a) *Aegilops* species, (b) *Avena sterilis*, (c) *Bromus sterilis*, (d) *Lolium rigidum*, (e) *Hordeum* species, and (f) *Hordeum* species richness.




FIGURE 2 (Continued)




FIGURE 2 (Continued)

M. rigidula (L.) Desr. and finally with *M. radiata* L., *M. sativa* L., and *M. minima* (L.) Bart. displaying lowest occurrence values. Species distribution map shows that *M. sativa* L. was only registered in Ain Ata-El Berke. Two major locations for high *Medicago* species richness were found in Al Fakiia and Sham El Hafour sites with four to five species (Figure 3d).

4.5.3 | *Trifolium*

Species distribution map shows its occurrence in all surveyed areas except on the very high elevations of Oro-Mediterranean shrubland, rocks, and screes habitat (Figure 3e). Seventeen species of *Trifolium* were identified. Four sites are designated as significant areas for these




FIGURE 3 Distribution map of (a) *Lathyrus* species, (b) *Lathyrus* species richness, (c) *Medicago* species, (d) *Medicago* species richness, (e) *Trifolium* species, and (f) *Trifolium* species richness in Mount Hermon.




FIGURE 3 (Continued)

species, most importantly Al Fakiaa and Sham El Hafour where the following 14 species were found: *Trifolium boisseri* Guss., *T. bullatum* var. *bullatum*, *T. campestre* Schreb., *T. clusii* Gren. & Godr. var. *clusii*, *T. fragiferum* L., *T. hirtum* All., *T. pilulare* Boiss., *T. purpureum* Boiss.,

T. resupinatum var. *resupinatum*, *T. scabrum* L., *T. scutatum* Boiss., *T. stellatum* L., *T. subterraneum* var. *brachycalycinum* Katzn. & Morley, and *T. tomentosum* var. *glabrescens* (Post) Hausskn. & Bornm. Another major area is Ain Ata- El Berke counting nine species of *Trifolium*.




FIGURE 3 (Continued)

Also, Rashaya Al Wadi-Wadi Sammouk contained species not found in the previous two areas, such as *T. arvense* L. and *T. bullatum* Boiss. & Hausskn var. *bullatum*. Species richness maps also show sites where a number up to seven species of *Trifolium* was concentrated (Figure 3f).

4.5.4 | *Vicia*

Vicia sativa L. and *V. tenuifolia* Roth. were identified in a limited distribution mainly in woodlands of Sham El Hafour (Figure 4a). Richness




FIGURE 4 Distribution maps of (a) *Vicia* species, (b) *Vicia* species richness, (c) *Lens culinaris* subsp. *orientalis* (Boiss.) Ponert, and (d) *Coronilla scorpioides* L. and *Pisum fulvum* Sibth. & Sm in Mount Hermon.

map shows that both species are found at the same time in only one location (Figure 4b).

Data collected for *Lens*, *Coronilla*, and *Pisum* are limited. *Lens* (Figure 4c) was mainly found in woodland-*Quercus infectoria* with

only one species (*Lens culinaris* subsp. *orientalis* (Boiss.) Ponert), especially in Al Yebse-Wadi El Botem (Figure 4d). *Coronilla scorpioides* L. also occurred primarily in woodland habitats, as well as *Pisum fulvum* Sibth. & Sm.




FIGURE 4 (Continued)

5 | DISCUSSION

This study reports the first comprehensive distribution mapping of a group of cereal and legume CWRs in Mount Hermon (Rachaya side) with a particular focus on genus and species richness and diversity

analyses being highly important in the protection and conservation of plant genetic resources for food and agriculture (Zair et al., 2020). The study highlights the importance of understanding CWR diversity and their distribution at both national and global levels. The species under study represent a substantial economic value to

agriculture and agrobiodiversity noting that endangered food crop relatives have a worth of about USD 10 billion annually in wholesale farm values (Phillips & Meilleur, 1998).

CWR genus and species richness maps (Figure 1a, b; Table 3) show that considering areas of highest richness values, such as Sham El Hafour Al Fakiaa and Ain Ata-al Berke, for a conservation plan could protect 40 out of the 46 targeted species. The remaining species either had low occurrences/frequency values in these two sites or their occurrence was prevalent in other study sites that are of specific habitat types. These findings are aligned with the study conducted by Bou Dagher Kharrat et al. (2018) that identified Mount Hermon as an important plant area due to its high richness index.

The analysis of sites with highest value of Shannon Index (Table 3) like Sham El Hafour (2.83) demonstrated the presence of 12 genera for 35 species, while each of Al Fakiaa (2.71), Ain Ata-al Berke (2.45), and Wadi El Raheb (2.41) sites retained nine genera designating, respectively 36, 24, and 21 species. Comparing the values of Shannon index found in Sham El Hafour and Al Fakiaa, where somewhat similar species richness values were found, higher densities of species were recorded in the former site. This result corroborates with findings of Akpo et al. (2000) that indicated a positive correlation between species richness and Shannon diversity index; the more numerous the species, the higher the Shannon-Weaver index. Thus, some of the species, at least, may dominate whatever are the conditions, considering that the ecological systems having more species should be steadier, and if a disruption occurs in the environment, the dominant species will be able to protect the whole community. The detailed map generated for Shannon index distribution shows that the values vary within the same habitat in the same location (Figure 1c). This could be attributed to the extent of various threats, mainly overgrazing, in the areas surveyed on the Mount. Other reasons could be accredited to the ethnobotanical aspect of the mountain where some species are collected seasonally by the local due to their benefits (Baydoun & Arnold, 2016).

The high occurrence of *Aegilops* species in 106 out of 170 surveyed quadrates shows a significant dominance for these species in the Mount (Table 2; Figure 2a). These findings are in line with previous unpublished surveys (in ICARDA's database) of *Aegilops* species in Mount Hermon by ICARDA researchers in 2020 resulting in the identification of *Ae. biuncialis*., *Ae. cylindrica*, *Ae. ovata*, *Ae. Triuncialis*, and *Ae. columnaris* in areas that were not included in our survey. This clearly indicates the high prevalence of these species in Mount Hermon. It is essential to mention that during the period when the field survey was achieved, a prevalent extent of the grazing threat was observed, which could have affected the identification and finding of other *Aegilops* species. Lebanon is diverse in *Aegilops*, as other studies reported the occurrence of nine *Aegilops* species in Nabha, and up to six species in Arsal and Ham (Amri et al., 2005). Highlighting the occurrence of this genus is essential due to its close relationship with cultivated wheat which has brought this genus subject to numerous studies (Bolot et al., 2009; Kumar et al., 2019; Xie et al., 2010). It represents a source of vast genetic diversity and

alleles of agronomic value to widen the wheat gene pool and improve tolerance to various types of potential stresses such pests, climate change, and other environmental stresses (Thiyagarajan et al., 2020). *Ae. biuncialis* Vis., specifically, is reported to be of a high potential for cross-pollination with the cultivated wheat cultivated bread wheat (*T. aestivum*) in Italy in optimized experimentation for hybridization (experimental soil layout, flowering synchrony) (Loureiro et al., 2006). Among other priority CWRs, this species is red-listed by IUCN meeting the category for Critically Endangered (CR) with high in situ priority (Perrino & Wagensommer, 2022). The flowering time of *Ae.* in Lebanon extends between April to June, depending on the species and its eco-geographical zone; this would suggest a high in situ crossbreeding potential.

On the contrary to *Aegilops*, one occurrence only of *Triticum* species (*T. aestivum*) was registered (Table 2). These results could be explained by the sampling method as ICARDA has previously reported records of *T. urartu* and *T. dicoccoides* in the Mount including the herein study sites knowing that these species had been first reported in the Mount by Mouterde (1951). According to the literature, these species having different and interesting genomic constitutions that can induce resistance to various environmental stresses, are among the main gene pool of domesticated wheat (Pour-Aboughadareh et al., 2021).

The presence of *A. sterilis* in all the habitats illustrated in Table 2 and Figure 2b aligns with Baum et al. (1972) stating the high adaptability of the genus considered a weed to a wide range of diverse habitats. A study of the phenotypic diversity of the species from Jordan revealed significant variations in different population traits illustrating the importance of considering various population clusters for gene enrichment in oat crop improvement (Al-Hajaj et al., 2018).

Similarly, *Lolium rigidum* Goud. was abundant in all habitat types. Actually, this species is considered by Terrell (1968) as invasive with high adaptability to different conditions. Researchers have identified *L. rigidum* and *L. multiflorum* as the species evolving glyphosate resistance in California based on morphology and assumed history (Jasieniuk et al., 2008; Simarmata et al., 2003).

As for *Bromus*, one species (*B. sterilis* L.) was only identified in the surveyed sites of this study (Table 2; Figure 2c), while occurrences of other species such as *B. alopecuroides*, *B. lanceolatus*, *B. tectorum*, and *B. tomentellus* were recorded in previous surveys conducted by Chalak et al. (n.d.) in similar habitats of the mountain.

With respect to *Lathyrus*, two species were identified in this study (Table 2; Figure 3a) as *L. aphaca* scored higher occurrence numbers and density than *L. blepharicarpus*. Currently 18 *Lathyrus* species are considered high priority species for conservation in the natural reserves of Lebanon like Horsh Ehdn National Reserve and Arz Bcharreh National Protected Zone (Shehadeh et al., 2013). This does not affect the possibility that this species could be also conserved in Mount Hermon as climatic conditions vary widely between the two established reserves and our study area. Different adaptability traits are suggested for the same *Lathyrus* species (Aci et al., 2011). In preliminary studies of interspecific hybridization with *L. sativus* Sibth. & Sm. ex Steud. (grasspea), some wild *Lathyrus* L. have

been found to be a potential germplasm resource in the breeding improvement programs of the grasspea (Kearney & Smartt, 1976; Yunus & Jackson, 1991).

In the case of *Medicago*, the species identified in our study had very different frequency and density values (Table 2) as well as a random selectivity of distribution in habitat types (Figure 3c). According to Beyrouthy et al. (2019), 38 species of medics are inventoried in Lebanon, of which 20 were found in protected areas including three species identified in Rachaya. In our study, we were able to collect accurate georeferenced occurrence data for eight *Medicago* species of high priority that could be added to the Lebanese distribution map for medics.

The distribution of the 17 identified *Trifolium* species was found in all the habitats and surveyed altitudes except in Fahet Jernaya-Beb El Hawa (2254 m a.s.l.) (Table 2, Figure 3e). Similar distribution trends were obtained in the study conducted by Atallah et al. (2008) as most of the *Trifolium* species appeared in all surveyed altitudes in the south bank of Nahr-Ibrahim valley. The absence of this species in the oro-Mediterranean shrublands, rocks, and screes habitat type is aligned with the survey conducted by (Chalak et al., n.d.) in Mount Hermon where none of the *Trifolium* species were found in these habitats. The two areas previously identified in this study as retaining the highest CWR diversity, Sham El Hafour and Al Fakiaa, comprise 14 out of the 17 identified species of *Trifolium*.

The lowest occurrence records for *V. sativa*, and *V. tenuifolia*, *Lens culinaris* subsp. *orientalis*, *Coronilla scorpioides*, *Pisum fulvum*, shown in Table 2 and Figure 4a, c, d may be attributed to the fact that these legume species are among the favorite for pasture grazing.

One of the important achievements of this study was that we were able to survey a new site (i.e., Chebaa-Ouyoun Jena) where no data was previously available. This site was the least explored region in the mountain due to security reasons. It lacked both historical and recent data (Bou Dagher Kharrat et al., 2018). In this site, we were able to identify 11 targeted species belonging to eight genera.

The Lebanese Parliament has voted to make Mount Hermon the country's 18th nature reserve. The designated protected area encompasses the sites we identified with highest species richness, Al Fakiaa, Sham El Hafour, and Ain Ata-al Berke. To effectively manage CWR species within this reserve, threat, habitat, population, and genetic assessments must be conducted to inform the development of a targeted management plan (Iriondo et al., 2008).

In situ conservation can be achieved through two strategies. The first strategy involves the implementation of conservation actions in areas where high diversity, high genus, and species richness of CWRs have been detected. This strategy targets the sites of Al Fakiaa, Sham El Hafour, and Ain Ata-al Berke. The second approach is at the genus level, a compilation of genus distribution maps and richness areas can be used to select conservation areas for projects focusing on specific genus or species. It is now an urgent priority to identify existing and novel mechanisms to finance and govern the global CWR network that will provide a fundamental basis for ensuring our future food security (Vincent et al., 2019).

6 | CONCLUSIONS

Recognizing the crucial importance of the protection of plant genetic resources, this study reports the first comprehensive mapping of CWRs distribution in Mount Hermon (Bekaa side) with the aim of selecting areas having highest priority species richness and with the perspective of recommending monitoring and conservation actions.

These findings play a crucial role in mapping CWRs in Lebanon, which is essential for enhancing both in situ and ex situ conservation of CWRs and agrobiodiversity, both nationally and internationally.

Based on diversity analysis and richness maps, we have identified three sites in Mount Hermon (i.e. Sham El Hafour, Al Fakiaa, and Ain Ata- al Berke) to be recommended for active CWR conservation in situ. These sites cover species that are primarily adapted to habitats of Woodland—*Quercus infectoria* and Agricultural land and Woodland—*Quercus coccifera*. The implementation of a conservation strategy in these sites has the potential to conserve 40 species out of the 46 identified CWR species belonging to 13 genera in Mount Hermon. Therefore, it is recommended, as a part of a management plan, to establish national policies that cover these sites, thus allowing stakeholders to become aware of the existence of a national landmark with global genetic importance. For the 6 missing CWR species from the recommended conservation sites, it's suggested to implement collecting efforts in order to make sure they are also conserved in at least ex situ manner.

Richness distribution maps produced in our study provide a clear idea of sites with highest species density for each genus for further collection missions aiming at complementing the in situ with ex situ conservation actions. Ex situ conservation is crucial and a gap analysis is necessary to determine which species and populations are yet to be conserved in genebanks. This analysis should encompass the broadest range of each species' genetic diversity to ensure active conservation. If a species is not currently conserved ex situ, collection missions can be arranged using the maps produced in this study to target various habitats and genetic material for breeding programs. If it is already conserved ex situ, an evaluation of the collection date is required to determine if new collection missions are necessary. Additionally, collection sites should also be chosen based on different habitats to preserve a wide range of genetic diversity for breeding purposes.

AUTHOR CONTRIBUTIONS

Eliane Sayde: Data curation (equal); formal analysis (equal); investigation (equal); project administration (equal); writing – original draft (equal). **Lamis Chalak:** Supervision (equal); writing – review and editing (equal). **Safaa Baydoun:** Project administration (supporting); resources (equal). **Ali Shehadeh:** Formal analysis (supporting); methodology (supporting); software (supporting). **Hicham El Zein:** Methodology (supporting); resources (supporting). **Jostelle Al Beyrouthy:** Project administration (supporting). **Mariana Yazbek:** Conceptualization (lead); funding acquisition (lead); methodology

(lead); resources (lead); supervision (equal); writing – review and editing (equal).

ACKNOWLEDGMENTS

This work was supported by a grant from ICARDA. Sincere appreciation goes to Miss Celine Sayde for her generous and fruitful assistance and to Mr. Mehdi Fayek, cultural and landscape guide in Mt. Hermon-Rachaya side for his valuable support.

CONFLICT OF INTEREST STATEMENT

The authors have no conflicts of interest to declare.

DATA AVAILABILITY STATEMENT

The data that supports the findings of this study are available in the supplementary material of this article.

ORCID

Hicham El Zein  <https://orcid.org/0000-0001-7541-9509>

Jostelle Al Beyrouthy  <https://orcid.org/0000-0002-0082-0936>

Mariana Yazbek  <https://orcid.org/0000-0001-8610-3809>

REFERENCES

- Aci, M. M., Lupini, A., Badagliacca, G., Mauceri, A., Lo Presti, E., & Preiti, G. (2020). Genetic diversity among *Lathyrus* ssp. based on agronomic traits and molecular markers. *Agronomy*, 10(8), 1182.
- Akpo, L. É., Grouzis, M., Bada, F., Pontanier, R., & Floret, C. (2000). Effet du couvert ligneux sur la structure de la végétation herbacée de jachères soudanaises. *Science et Changements Planétaires/ Sécheresse*, 10(4), 253–261.
- Al-Atawneh, N., Amri, A., Assi, R., & Maxted, N. (2008). Management plans for promoting in situ conservation of local agrobiodiversity in the West Asia centre of plant diversity. In N. Maxted, B. V. Ford-Lloyd, S. P. Kell, J. Iriondo, M. E. Dulloo, & K. Turuk (Eds.), *Crop wild relative conservation and use* (pp. 338–361). CABI Publishing.
- Al Beyrouthy, J., Karam, N., Al-Zein, M. S., & Yazbek, M. (2019). Ecogeographic survey and gap analysis for *Medicago* L.: Recommendations for in situ and ex situ conservation of Lebanese species. *Genetic Resources and Crop Evolution*, 66(5), 1009–1026.
- Al-Hajaj, N., Peterson, G. W., Horbach, C., Al-Shamaa, K., Tinker, N. A., & Fu, Y. B. (2018). Genotyping-by-sequencing empowered genetic diversity analysis of Jordanian oat wild relative *Avena sterilis*. *Genetic Resources and Crop Evolution*, 65, 2069–2082.
- Amri, A., Ajlouni, M., Sbeih, Y., Assi, R., & Khnifes, A. (2005). Promoting the conservation and sustainable use of dryland agrobiodiversity. In the Proceeding of International Conference on “Promoting community-driven in situ conservation of dryland agrobiodiversity” organized 19–25 April 2005.
- Amri, A., Yazbek, M., Shehadeh, A., Nawar, M., Tsivelikas, A., & Biradar, C. (2016). ICARDA efforts to promote in situ / on-farm conservation of dryland agrobiodiversity. *Indian Journal of Plant Genetic Resources*, 29, 265. <https://doi.org/10.5958/0976-1926.2016.00042.5>
- Arnold, N., Baydoun, S., Chalak, L., & Raus, T. (2015). A contribution to the flora and ethnobotanical knowledge of mount Hermon, Lebanon. *Flora Mediterranea*, 25, 13–55.
- Atallah, T., Rizk, H., Cherfane, A., Daher, F. B., El-Alia, R., De Lajuide, P., & Hajj, S. (2008). Distribution and nodulation of spontaneous legume species in grasslands and shrublands in Mediterranean Lebanon. *Arid Land Research and Management*, 22(2), 109–122.
- Baum, B. R., Fleischmann, G., Martens, J. W., Rajhathy, T., & Thomas, H. (1972). Notes on the habitat and distribution of *Avena* species in the Mediterranean and Middle East. *Canadian Journal of Botany*, 50(6), 1385–1397.
- Baydoun, S., & Arnold, N. (2016). Ethnobotanical study of wild edible plants in rural communities of Lebanon. Abstract and poster presented at XV OPTIMA Meeting, Montpellier (France).
- Baydoun, S., Chalak, L., Dalleh, H., & Arnold, N. (2015). Ethnopharmacological survey of medicinal plants used in traditional medicine by the communities of mount Hermon, Lebanon. *Journal of Ethnopharmacology*, 173, 139–156.
- Bierschenk, B., Tagele, M. T., Ali, B., Ashrafuzzaman, M. D., Wu, L. B., Becker, M., & Frei, M. (2020). Evaluation of rice wild relatives as a source of traits for adaptation to iron toxicity and enhanced grain quality. *PLoS ONE*, 15(1), e0223086.
- Bohra, A., Kilian, B., Sivasankar, S., Caccamo, M., Mba, C., McCouch, S. R., & Varshney, R. K. (2021). Reap the crop wild relatives for breeding future crops. *Trends in Biotechnology*, 40(4), 412–431.
- Bolot, S., Abrouk, M., Masood-Quraishi, U., Stein, N., Messing, J., Feuillet, C., & Salse, J. (2009). The ‘inner circle’ of the cereal genomes. *Current Opinion in Plant Biology*, 12(2), 119–125.
- Bou Dagher Kharrat, M., El Zein, H., & Rouhan, G. (2018). Setting conservation priorities for Lebanese flora—Identification of important plant areas.
- Castañeda-Álvarez, N., Khoury, C., Achicanoy, H., Guarino, L., Harker, R. H., Jarvis, A., Maxted, N., Müller, J. V., Ramirez-Villegas, J., Sosa, C. C., Struik, P. C., Vincent, H., & Toll, J. (2016). Global conservation priorities for crop wild relatives. *Nature Plants*, 2, 16022. <https://doi.org/10.1038/nplants.2016.22>
- Ceccarelli, S., Maxted, N., Dulloo, M. E., Ford-Lloyd, B. V., Frese, L., Iriondo, J. M., & de Carvalho, M. P. (2012). Agrobiodiversity conservation: Securing the diversity of crop wild relatives and landraces.
- Chalak, L., El Zein, H., Kkater, C., Yazbek, M., & Baydoun, S. (n.d.). Assessing and mapping habitats of Hermon Mountain nature reserve. Submitted for publication.
- Chalak, L., Mba, C., Diulgheroff, S., & Muhammad, D. (2020). Development of National Strategies for the conservation and sustainable utilization of plant genetic resources for food and agriculture in some countries of the near east and North Africa. *Acta Horticulturae*, 1297, 9–16.
- Chalak, L., Noun, J., El Haj, S., Rizk, H., Assi, R., Attieh, J., Maalouf, F., Abi Antoun, M., & Sabra, N. (2011). Current status of agro-biodiversity in Lebanon and future challenges. *Gene Conserve*, 10(39), 23–41.
- Euro Med. (2006). Euro+Med Plant Base—the information resource for Euro-Mediterranean plant diversity.
- Gaston, K. J. (1996). Spatial patterns in taxonomic diversity. In *Biodiversity: A biology of numbers and difference* (pp. 77–113). Wiley-Blackwell.
- GBIF.org. (2021). GBIF Home Page. <https://www.gbif.org>
- Good, I. J. (1953). The population frequencies of species and the estimation of population parameters. *Biometrika*, 40(3-4), 237–264.
- Iriondo, J. M., Maxted, N., & Dulloo, M. E. (2008). *Conserving plant genetic diversity in protected areas: Population management of crop wild relatives*. CABI.
- IUCN. (2001). *IUCN red list categories and criteria: Version 3.1. IUCN species survival commission*. IUCN.
- Jarvis, A., Lane, A., & Hijmans, R. J. (2008). The effect of climate change on crop wild relatives. *Agriculture, Ecosystems & Environment*, 126(1-2), 13–23.
- Jasieniuk, M., Ahmad, R., Sherwood, A. M., Firestone, J. L., Perez-Jones, A., Lanini, W. T., Mallory-Smith, C., & Stednick, Z. (2008). Glyphosate-resistant Italian ryegrass (*Lolium multiflorum*) in California: distribution, response to glyphosate, and molecular evidence for an altered target enzyme. *Weed Science*, 56(4), 496–502.
- Jenks, G. F. (1967). The data model concept in statistical mapping. *International Yearbook of Cartography*, 7, 186–190.
- Kearney, J., & Smartt, J. (1976). The grasspea *Lathyrus sativus* (Leguminosae-Papilionoideae). In N. W. Simmonds (Ed.), *In evolution of crop plants*. Longman.

- Kell, S. P., Maxted, N., & Bilz, M. (2012). European crop wild relative threat assessment: Knowledge gained and lessons learnt. In N. Maxted, M. E. Dulloo, B. V. Ford-Lloyd, L. Frese, J. M. Iriondo, & M. A. A. Pinheiro de Carvalho (Eds.), *Agrobiodiversity conservation: Securing the diversity of crop wild relatives and landraces*. CAB International.
- Khoury, C. K., Argumedo, A., Dempewolf, H., Guarino, L., Haan, S. D., Wenzl, P., DeClerk, F., Dulloo, E., Hunter, D., Padulosi, S., & Etten, J. V. (2019). *With Diversity at the heart: a 2050 Vision for Global Food System Transformation*.
- Kumar, A., Kapoor, P., Chunduri, V., Sharma, S., & Garg, M. (2019). Potential of *Aegilops* sp. for improvement of grain processing and nutritional quality in wheat (*Triticum aestivum*). *Frontiers in Plant Science*, 10, 308.
- Loureiro, I., Escorial, M. C., García-Baudin, J. M., & Chueca, M. C. (2006). Hybridization between wheat (*Triticum aestivum*) and the wild species *Aegilops geniculata* and *A. biuncialis* under experimental field conditions. *Agriculture, Ecosystems and Environment*, 120, 384–390.
- MacKenzie, D. I., Nichols, J. D., Royale, J. A., Pollock, K. H., Bailey, L. L., & Hines, J. E. (2018). *Occupancy estimation and modeling: Inferring patterns of dynamics of species occurrence* (2nd ed.). Academic Press.
- Magurran, A. E. (Ed.). (1988). *Diversity indices and species abundance models. Ecological diversity and its measurement*. Springer. https://doi.org/10.1007/978-94-015-7358-0_2
- Maxted, N., Amri, A., Castañeda-Álvarez, N. P., Dias, S., Dulloo, M. E., Fielder, H., Ford-Lloyd, B. V., Iriondo, J. M., Brehm, J. M., Nilsen, L. B., Thormann, I., Vincent, H., & Kell, S. P. (2016). Joining up the dots: a systematic perspective of crop wild relative conservation and use. In *Enhancing crop gene pool use: capturing wild relative and landrace diversity for crop improvement* (pp. 87–124). CABI.
- Medail, F., & Quezel, P. (1997). Hot-spots analysis for conservation of plant biodiversity in the Mediterranean Basin. *Annals of the Missouri Botanical Garden*, 84, 112–127.
- Mouterde, P. (1966, 1970, 1983). *New Flora of Lebanon and Syria Beirut*. Catholic Printing.
- Mouterde, P. (1986). *Nouvelle flore du Liban et de la Syrie, vol 2*. Dar el-Machreq Editeurs.
- Mouterde, R. (1951). *Antiquités de l'Hermon et de la Beqâ*. Mélanges de l'Université Saint-Joseph.
- Nail, K. P. (2019). Utilizing crop wild relatives to combat global warming. *Advances in Agronomy*, 153, 175–258.
- Ng'uni, D., Munkombwe, G., Mwila, G., Gaisberger, H., Brehm, J. M., Maxted, N., Kell, S., & Thormann, I. (2019). Spatial analyses of occurrence data of crop wild relatives (CWR) taxa as tools for selection of sites for conservation of priority CWR in Zambia. *Plant Genetic Resources*, 17(2), 103–114.
- Pacicco, L., Bodesmo, M., Torricelli, R., & Negri, V. (2018). A methodological approach to identify agro-biodiversity hotspots for priority in situ conservation of plant genetic resources. *PLoS ONE*, 13(6), e0197709.
- Perrino, E. V., & Wagensommer, R. P. (2021). Crop wild relatives (CWR) priority in Italy: Distribution, ecology, in situ and ex situ conservation and expected actions. *Sustainability*, 13(4), 1682.
- Perrino, E. V., & Wagensommer, R. P. (2022). Crop wild relatives (CWRs) threatened and endemic to Italy: Urgent actions for protection and use. *Biology*, 11(2), 193.
- Phillips, O. L., & Meilleur, B. (1998). Usefulness and economic potential of the rare plants of the United States: A status survey. *Economic Botany*, 52, 57–67.
- Pour-Aboughadareh, A., Kianersi, F., Poczai, P., & Moradkhani, H. (2021). Potential of wild relatives of wheat: Ideal genetic resources for future breeding programs. *Agronomy*, 11, 1656. <https://doi.org/10.3390/agronomy11081656>
- Rajan, S. S. (2001). *Practical manual of plant ecology and plant physiology* (1st ed.). ANMOL, Publications. PVT.LTD.
- Ratnayake, S. S., Kariyawasam, C. S., Kumar, L., Hunter, D., & Liyanage, A. S. U. (2021). Potential distribution of crop wild relatives under climate change in Sri Lanka: Implications for conservation of agricultural biodiversity. *Current Research in Environmental Sustainability*, 3, 100092.
- Shehadeh, A., Amri, A., & Maxted, N. (2013). Ecogeographic survey and gap analysis of lathyrus L. species. *Genetic Resources and Crop Evolution*, 60(7), 2101–2113.
- Simarmata, M., Kaufmann, J. E., & Penner, D. (2003). Potential basis of glyphosate resistance in California rigid ryegrass (*Lolium rigidum*). *Weed Science*, 51(5), 678–682.
- Terrell, E. E. (1968). A taxonomic revision of the genus *Lolium*. (No. 1392). Agricultural Research Service. US Department of Agriculture.
- Thiyagarajan, K., Latini, A., Cantale, C., & Galeffi, P. (2020). Structural characterization of the DRF1 gene of *Ae. speltooides* and comparison of its sequence with those of B and other *Triticeae* genomes. *Euphytica*, 216, 152.
- Vincent, H., Amri, A., Castañeda-Álvarez, N. P., Dempewolf, H., Dulloo, E., Guarino, L., Guarino, L., Hole, D., Mba, C., Toledo, A., & Maxted, N. (2019). Modeling of crop wild relative species identifies areas globally for in situ conservation. *Communications Biology*, 2(1), 136.
- Vincent, H., Wiersema, J., Kell, S., Fielder, H., Dobbie, S., Castañeda-Álvarez, N. P., Guarino, L., Eastwood, R., León, B., & Maxted, N. (2013). A prioritized crop wild relative inventory to help underpin global food security. *Biological Conservation*, 167, 265–275.
- Xie, Z., Wang, C., Wang, K., Wang, S., Li, X., Zhang, Z., Ma, W., & Yan, Y. (2010). Molecular characterization of the celiac disease epitope domains in α -gliadin genes in *Ae. tauschii* and hexaploid wheats (*Triticum aestivum* L.). *Theoretical and Applied Genetics*, 121(7), 1239–1251.
- Yunus, A. G., & Jackson, M. T. (1991). The gene pools of the grasspea (*Lathyrus sativus* L.). *Plant Breeding*, 106, 319–328.
- Zair, W., Maxted, N., & Amri, A. (2018). Setting conservation priorities for crop wild relatives in the Fertile Crescent. *Genetic Resources and Crop Evolution*, 65, 855–863.
- Zair, W., Maxted, N., Brehm, J. M., & Amri, A. (2020). Ex situ and in situ conservation gap analysis of crop wild relative diversity in the Fertile Crescent of the Middle East. *Genetic Resources and Crop Evolution*, 68, 693–709.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Sayde, E., Chalak, L., Baydoun, S., Shehadeh, A., El Zein, H., Al Beyrouthy, J., & Yazbek, M. (2024). Surveying and mapping cereals and legumes wild relatives in Mount Hermon (Bekaa, Lebanon). *Ecology and Evolution*, 14, e10943. <https://doi.org/10.1002/ece3.10943>